

LA-UR-18-28474

Approved for public release; distribution is unlimited.

Title: High-energy Particle Physics -- In Space!


Author(s): Albert, Andrea

Intended for: LANL NEN Seminar


Issued: 2018-09-05

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.



Fermi LAT & HAWC Collaborations



High-energy Particle Physics In Space!

Andrea Albert
Los Alamos National Laboratory

NEN Seminar
September 6, 2018

Photons are an Astronomical Probe




Photons as an Astronomical Probe




**Quiver Tree Forest
Namibia Africa**

<http://apod.nasa.gov/apod/ap121212.html>

Different Wavelengths Probe Different Physics



Gamma rays Probe the Extreme, Non-Thermal, Universe



Gamma rays Probe the Extreme, Non-Thermal, Universe


Dark Nebula




Dim, young star



Our Sun




Globular Cluster



Visible photons $E \sim 1 \text{ eV}$

Gamma-ray photons $E \sim \text{GeV} (1,000,000,000 \text{ eV})$




Non-thermal γ -ray Emission



Energy source



Acceleration
mechanism



γ -ray production
mechanism






Foreground absorption




γ rays

Emission



Inverse Compton (IC)





dark matter annihilation???







γ -ray production mechanism

γ -ray production mechanisms are similar to interactions investigated by high-energy particle physics experiments like those using the Large Hadron Collider

Different Wavelengths Require Different Telescope Technologies



Different Wavelengths Require Different Telescope Technologies




Current Gamma-ray Observatories




Tucson, Arizona
31° North Latitude, ~5° f.o.v.
~85 GeV -- ~50 TeV




La Palma, Canary Islands
29° North Latitude, ~5° f.o.v.
~30 GeV -- ~30 TeV



Khomas Highland of Namibia
23° South Latitude, ~5° f.o.v.
~30 GeV -- ~100 TeV



Parque Nacional Pico de Orizaba, Mexico
19° North Latitude, ~2 sr f.o.v.
~50 GeV -- ~100 TeV, ~100% Duty Cycle



Low earth orbit (565 km)
28.5° orbital inclination, ~2 sr f.o.v.
20 MeV -- > 300 GeV, ~100% Duty Cycle
(AGILE has similar technology, but has limited energy resolution)


Current Gamma-ray Observatories



Tucson, Arizona
31° North Latitude, ~5° f.o.v.
~85 GeV -- ~50 TeV



La Palma, Canary Islands
29° North Latitude, ~5° f.o.v.
~30 GeV -- ~30 TeV



Khomas Highland of Namibia
23° South Latitude, ~5° f.o.v.
~30 GeV -- ~100 TeV

Air Cherenkov Telescopes (ACTs) have deeper exposure, better angular resolution, but smaller field of view and only operate at night in during good weather


Pa
19° North Latitude, ~2° f.o.v.
~50 GeV -- ~100 TeV, ~100% Duty Cycle

(limited energy resolution)

HAWC (~1 TeV) and Fermi LAT (~1 GeV) are complementary wide field of view survey instruments. Provide continuous monitoring of the gamma-ray sky across 7 decades of energy (20 MeV to >100 TeV)



HAWC Observatory
Parque Nacional Pico de Orizaba, Mexico
19° North Latitude, ~2 sr f.o.v.
~50 GeV -- ~100 TeV, ~100% Duty Cycle




Fermi Large Area Telescope
Low earth orbit (565 km)
28.5° orbital inclination, ~2 sr f.o.v.
20 MeV -- > 300 GeV, ~100% Duty Cycle
(AGILE has similar technology, but has limited energy resolution)

- On board the Fermi Gamma-ray Space Telescope
 - Launched June 11, 2008 (science mission started Aug. 2008)
 - Mission extended at least through 2018
 - No consumables
 - Orbit re-entry expected ~2026-2044 (depending on solar activity)
 - LAT has triggered on ~500 billion events
 - Processed ~100 billion events (>1 Petabyte!)

Large Area Telescope (LAT)

Observes 20% of the sky at any instant, views entire sky every 3 hrs
20 MeV - 300 GeV - includes unexplored region between 10 - 100 GeV

Can go >300 GeV



Fermi Large Area Telescope (LAT)


- On 11 Feb 2008, the Fermi Gamma-ray Space Telescope was launched (Aug. 2008)

Large
Observ
20 MeV

Can go
my Ph.D. adviser Dr. Brian Winer got
his facebook profile pic at launch



on solar activity)



Gamma-ray Burst Monitor (GBM)
Observes entire unocculted sky
Detects transients from 8 keV - 40 MeV





Public Data Release:

All γ -ray data made public
within 24 hours (usually less)

Si-Strip Tracker:

convert $\gamma \rightarrow e^+e^-$
reconstruct γ direction
EM v. hadron separation

Hodoscopic CsI Calorimeter:


measure γ energy
image EM shower
EM v. hadron separation

Trigger and Filter:

Reduce data rate from $\sim 10\text{kHz}$
to 300-500 Hz

Fermi LAT Collaboration:




~ 400 Scientific Members,
NASA / DOE & International
Contributions



Simulated 27 GeV Gamma-ray Event





Incoming γ -ray direction






Simulated 27 GeV Gamma-ray Event



Incoming γ -ray direction




Simulated 27 GeV Gamma-ray Event




Simulated 27 GeV Gamma-ray Event

ACD hits from
backsplash



**Hodoscopic CAL gives
provides 3D directional info**

LAT Simulated 1 TeV Gamma-ray Event



The HAWC Observatory

Joint NSF/DOE/CONACyT (Mexico)

HAWC Observatory

HAWC operates day and night, providing a large field of view for the observation of the highest energy gamma rays.



~17 rad
lengths


Pico de Orizaba
(5,626 m)

HAWC is located at 4,100 m
above sea level, covering
an area of 20,000 m².

Andrea Albert (LANL)

Water Cherenkov tank


HAWC comprises an array of 300 tanks that record the particles created in gamma-ray and cosmic-ray showers.



Gamma rays vs cosmic rays


HAWC selects gamma rays from among a much more abundant background of cosmic rays.

gamma-ray shower



"hot" spots concentrate
around the core

cosmic-ray shower



"hot" spots are more
dispersed

The HAWC Observatory


Joint NSF/DOE/CONACyT (Mexico)

current supervisor's facebook profile pic (Dr. Brenda Dingus)



Water Cherenkov tank


HAWC comprises an array of 300 tanks that record the particles created in gamma-ray and cosmic-ray showers.



My colleague Kelly Malone (graduate student at Penn State)





HAWC is located at 4,100 m above sea level, covering an area of 20,000 m².



Andrea Alber

The HAWC Observatory


Water Cherenkov tank



Gamma rays vs cosmic rays


HAWC selects gamma rays from among a much more abundant background of cosmic rays.

gamma-ray shower




"hot" spots concentrate around the core

cosmic-ray shower





"hot" spots are more dispersed

HAWC is located at 4,100 m above sea level, covering an area of 20,000 m².





~60 TeV gamma ray
event from the Crab




**Angle of primary is
determined from
timing.**



**Energy of primary is
determined from
energy in detector.**









Fermi Large Area Telescope 7 year map >1 GeV




Fermi LAT GeV Gamma-ray Sky




**Markarian 501
(AGN)** →







Fermi LAT GeV Gamma-ray Sky






Fermi LAT GeV Gamma-ray Sky








HAWC 1 year >1 TeV



HAWC 1 year >1 TeV




HAWC 1 year >1 TeV Equatorial Coordinates



At a latitude of $\sim 20^\circ$, HAWC can see the sky from about Declinations $+60^\circ$ to -20°

What is the source of high-energy cosmic rays?



- Here 'cosmic rays' = particles like protons and electrons
 - different than γ rays


We observe particles with 10,000,000 times more energy than the collisions at the LHC

- Where do these come from?
- What is giving them all that energy?

Highest energy events are rare and require huge arrays like Pierre Auger

- Pierre Auger area is $\sim 1,000 \text{ km}^2$
- 'Oh my god' particle observed with $E = 3 \times 10^{20} \text{ eV}$ in 1991 with Fly's Eye (Utah)

Cosmic-ray Protons should make pions





- high-energy protons will create pions
 - neutral pions quickly decay to a pair of gamma rays
 - energy spectrum of gamma rays from pion decay would peak at $m_{\pi^0}/2 \sim 70 \text{ MeV}$
- Supernova remnants are possible sources of cosmic-ray proton acceleration
 - to test this hypothesis, we can search for the 'pion bump' using the Fermi LAT




magenta = Fermi GeV gamma-ray emission,
yellow = optical wavelengths,
blue, cyan, green, red = infrared data (3.4, 4.6, 12, 22 microns)

Credit: NASA/DOE/Fermi LAT Collaboration,
NOAO/AURA/NSF, JPL-Caltech/UCLA
https://www.nasa.gov/mission_pages/GLAST/news/supernova-cosmic-rays.html

Pion Bump Seen in Supernova Remnants







- pion bump seen from two supernova remnants: IC 443, W44
 - direct detection that supernova remnants are cosmic-ray proton accelerators
- protons required to make these gamma rays only require energies up to ~1 TeV
 - doesn't explain 'oh my god' particle



Fermi-LAT gamma-ray image of two supernova remnants: W44 and IC 443


The Crab Nebula





Crab Nebula in various wavelengths
credit: GFSC/NASA

- Space must also accelerate electrons
 - high energy electrons observed by e.g. AMS
- Know Pulsar Wind Nebula power electrons since we see synchrotron emission in radio/microwave
 - high energy electrons upscatter Cosmic Microwave Background (CMB) photons to gamma rays
 - inverse Compton (IC)

Andrea Albert (LANL)





The Crab Nebula -- Looking for the Highest Energy Gamma Ray



- Inverse Compton component decreases rapidly at high energies
 - electrons lose energy too quickly to get energetic enough to create greater than about 60 TeV gamma rays
 - high energy protons are expected to create highest energy gamma rays
- HAWC is primed to observe the highest energy gamma ray ever detected
 - upgrade underway to expand array and improve sensitivity above 10 TeV by a factor of 3-4



Upgrade to HAWC Array

A. Sandolval arXiv:1509.04269




- Upgrade to HAWC array is underway
 - add larger, sparse array of small tanks
- Provide better measure of high energy showers
 - expect gain in sensitivity > 10 TeV of about 3-4

- Dark Matter is the **gravitational glue** that holds e.g. galaxies together



- Weakly Interacting Massive Particle
 - promising DM candidate
 - fundamental particle beyond Standard Model
 - Being searched for at LHC and underground experiments (e.g. LUX)
 - WIMP annihilations may produce γ rays**
 - weak scale annihilation cross section \rightarrow observed abundance of DM




- Dark Matter is the **gravitational glue** that

Gamma-rays




neutrinos

d
er




- Weakly
 - pro
 - fun
 - WI


Milky Way Dwarf Spheroidal Galaxies



Milky Way Dwarf Spheroidal Galaxies




Gamma Rays from Dark Matter




σ is the annihilation cross section. Larger σ means there is a larger chance of interaction.

Limits from Dwarf Galaxies



- No gamma-ray excess observed in dwarf galaxies
- Together Fermi LAT and HAWC limits constrain dark matter models for masses from ~2 GeV to 100 TeV

Limits from Dwarf Galaxies



- No gamma-ray excess observed in dwarf galaxies
- Together Fermi LAT and HAWC limits constrain dark matter models for masses from ~ 2 GeV to 100 TeV

- The Fermi LAT is an **awesome** particle detector in space!
 - in orbit for 8 years and counting
- HAWC is an **awesome** particle detector in the highest mountains in Mexico!
 - Operating with full array since March 2015
 - New detector expansion underway to reach highest gamma ray energies
- Gamma rays tell us about high energy particle physical mechanisms in space
 - help us answer fundamental questions
 - “where/how are the high energy cosmic rays accelerated”
 - “what are the particle properties of dark matter”

Backup




- **Weakly Interacting Massive Particle (WIMP, χ)**

- GeV - TeV mass scale
- WIMPs may be thermal relics
- e.g. neutralino (SUSY, electrically neutral, stable, motivated theoretically)


Assuming a weak scale σ_{ann} yields observed relic abundance

- $\langle \sigma v \rangle_{\text{ann}} \sim 3 \times 10^{-26} \text{ cm}^3/\text{s}$ ($\sigma_{\text{ann}} \sim 3 \text{ pb}$ at time of decoupling)


Particle DM annihilation may produce Standard Model particles



Indirect Dark Matter Detection



Broad spectrum



- The Milky Way sits in a large halo of cold DM ($v_{DM} \ll c$)
 - Expect additional DM overdensities (halos / subhalos)
 - e.g. Milky Way dwarf galaxies
 - e.g. Galaxy Clusters

WIMP annihilations (decays) may produce gamma rays

